Color s:



bait containing Cholecalciferol 0.077 % w/w (0.075 % w/w pure). AUTHORISED USES:

GB. NI and IRFI AND

Harmonix Rodent Paste controls:

- . House Mice (Mus musculus)
- . Norway Rats (Rattus norvegicus) · Black Rats (Rattus rattus)



For trained professional users

For professional users with demonstrated competence (equivalent to trained professionals)

: Indoor use, outdoor (around buildings) and outdoor (open areas and waste dumps). APPLICATION METHOD:





Baiting for Mice: 20 g (1 sachet) bait per baiting point depending upon infestation level. The minimum distance between baiting points should be 5-20 linear metres (in case of a low infestation) and 2-5 linear metres (in case of a high infestation).

Refill an amount of 20 g (1 sachet) to a bait point if bait has been totally consumed within one control

Baiting for Rats: 100-200 g (5-10 sachets) bait per baiting point depending upon infestation level. The minimum distance between baiting points should be 10-20 linear metres (in case of a low infestation) and 3-10 linear metres (in case of a high infestation). Refill an amount of 200 g (10 sachets) to a bait point if bait has been totally consumed within one control interval.





Indoor use, outdoor (around buildings) and outdoor (open areas and waste dumps).

Target organism	Dose (per baiting point)	Level of infestation	Distance between baiting points – tamper resistant bait stations or covered and protected baiting points
Mice	20 g (1 sachet)	High	2-5 linear metres
		Low	5-20 linear metres
Rats	100-200 g (5-10 sachets)	High	3-10 linear metres
		Low	10-20 linear metres





GB. NI and IRFLAND

For trained professional users. For professional users with demonstrated competence (equivalent to trained professionals)

Direct application of ready-to-use bait into the burrow (outdoor around buildings and outdoor open areas and waste dumps only): Baits must be placed to minimise the exposure to non-target species and children. Cover or block the entrances of baited burrows to reduce the risks of bait being rejected and spilled.

Permanent baiting: should be done around nesting sites, entry points or immediate vicinity of buildings using the same amount of bait per baiting point. Sites under a permanent baiting regime should be inspected regularly. The period between visits should be determined by the technician in charge but will not be longer than every four weeks when permanent baiting is conducted outdoors. For permanent baiting follow any additional instructions provided by the CRRU Guidance on Permanent Baiting.

