

## GB - RATIMOR BRODIFACUM FRESH BAIT

### INSTRUCTIONS FOR USE:

#### Indoor (House mouse, Brown rat and Black or Roof rat):

For mouse infestations use 10 g up to 20 g of bait per bait point. Place bait points 5 m apart reducing to 2 m in high infestations. The bait stations should be visited at least every 2 to 3 days at the beginning of the treatment and at least weekly afterwards, in order to check whether the bait is accepted, the bait stations are intact and to remove rodent bodies. Re-fill bait when necessary. For rat infestations use 10 g up to 60 g of bait per bait points. Place bait points 10 m apart reducing to 5 m in high infestations. The bait stations should be visited only 5 to 7 days after the beginning of the treatment and at least weekly afterwards, in order to check whether the bait is accepted, the bait stations are intact and to remove rodent bodies. Re-fill bait when necessary. Ready-to-use bait to be used in tamper-resistant bait stations. Ready-to-use bait to be used in covered and protected baiting points.

Remove the remaining product at the end of treatment period. If used for pulsed baiting: Replace eaten bait only after 3 days and than at maximum 7 day intervals. Collect any spilled bait and dead rodents.

Where possible, prior to the treatment inform any possible bystanders (e.g. users of the treated area and their surroundings) about the rodent control campaign. Consider preventive control measures (e.g. plug holes, remove potential food and drinking as far as possible) to improve product intake and reduce the likelihood of reinfestation. To reduce risk of secondary poisoning, search for and remove dead rodents during treatment at frequent intervals, in line with the recommendations provided by the relevant code of best practice. Do not use the product as permanent baits for the prevention of rodent infestation or monitoring of rodent activities. When placing bait points close to water drainage systems, ensure that bait contact with water is avoided.

#### Outdoor around buildings (House mouse, Brown rat and Black or Roof rat):

For mouse infestations use 10 g up to 20 g of bait per bait point. Place bait points 5 m apart reducing to 2 m in high infestations. The bait stations should be visited at least every 2 to 3 days at the beginning of the treatment and at least weekly afterwards, in order to check whether the bait is accepted, the bait stations are intact and to remove rodent bodies. Re-fill bait when necessary. For rat infestations use 10 g up to 60 g of bait per bait points. Place bait points 10 m apart reducing to 5 m in high infestations. The bait stations should be visited only 5 to 7 days after the beginning of the treatment and at least weekly afterwards, in order to check whether the bait is accepted, the bait stations are intact and to remove rodent bodies. Re-fill bait when necessary. Ready-to-use bait to be used in tamper-resistant bait stations. Ready-to-use bait to be used in covered and protected baiting points. Direct application of ready-to-use bait into the burrow.

Protect bait from the atmospheric conditions. Place the baiting points in areas not liable to flooding. Replace any bait in baiting points in which bait has been damaged by water or contaminated by dirt. For outdoor use, baiting points must be covered and placed in strategic sites to minimise the exposure to non-target species. Remove the remaining product at the end of treatment period (except when directly applied into burrows). If used for pulsed baiting: Replace eaten bait only after 3 days and than at maximum 7 day intervals. Collect any spilled bait and dead rodents. When used in burrows: Baits must be placed to minimise the exposure to non-target species and children. Cover or block the entrances of baited burrows to reduce the risks of bait being rejected and spilled. Follow any additional instructions provided by the CRRU UK Code of Best Practice (or equivalent).

Where possible, prior to the treatment inform any possible bystanders (e.g. users of the treated area and their surroundings) about the rodent control campaign. Consider preventive control measures (plug holes, remove potential food and drinking as far as possible) to improve product intake and reduce the likelihood of reinfestation. To reduce risk of secondary poisoning, search for and remove dead rodents during treatment at frequent intervals, in line with the recommendations provided by the relevant code of best practice. Do not use the product as permanent baits for the prevention of rodent infestation or monitoring of rodent activities. When placing bait points close to surface waters (e.g. rivers, ponds, water channels, dykes, irrigation ditches) or water drainage systems, ensure that bait contact with water is avoided.

## UNICHEM

**Manufacturer and approval holder:**  
UNICHEM d.o.o., Sinja Gorica 2, 1360 Vrhnika,  
Slovenia, Tel.: +386 1 7558 152,  
unicchem@unicchem.si, www.unicchem.si

#### Distributor:

Killgerr Chemicals Ltd, P.O. BOX 2, Ossett,  
West Yorkshire WFS 9NA, UK,  
Tel.: 01924 268 400, Fax: 01924 264 757,  
www.killgerr.com

# RATIMOR®

## BRODIFACUM

## 0,029 g/kg

### Fresh bait 60 g

### WITH SPECIAL AROMA



## 3 kg

#### Sewers (Brown rat):

For rat infestations in sewers use up to 200 g per manhole. In case of high infestations use 200 g per manhole.

Ready-to-use bait to be anchored or applied in bait stations preventing the bait from getting into contact with waste water. Covered and protected baiting points preventing the bait from getting into contact with waste water.

Baits must be applied in a way so that they do not come into contact with water and are not washed away.

Do not use the product in pulsed baiting treatments or as permanent baits for the prevention of rodent infestation or monitoring of rodent activities.

**FIRST AID:** This product contains an anticoagulant substance. If ingested, symptoms, which may be delayed, may include nosebleed and bleeding gums. In severe cases, there may be bruising and blood present in the faeces or urine. Antidote: Vitamin K1 administered by medical/veterinary personnel only. In case of: - Dermal exposure, wash skin with water and then with water and soap. - Eye exposure, rinse eyes with eyes-rinse liquid or water, keep eyes lids open at least 10 minutes. - Oral exposure, rinse mouth carefully with water. Never give anything by mouth to unconscious person. Do not provoke vomiting. If swallowed, seek medical advice immediately and show the product's container or label. UK medical professionals should contact the National Poisons Information Service ([www.npis.org](http://www.npis.org)) for further advice. Contact a veterinary surgeon in case of ingestion by a pet. **Bait stations must be labelled with the following information: "do not move or open"; "contains a rodenticide"; "product name or authorisation number"; "active substance(s)" and "in case of incident, call a poison centre".** Hazardous to wildlife.

Because of their delayed mode of action, anticoagulant rodenticides take from 4 to 10 days to be effective after consumption of the bait. Rodents can be disease carriers. Do not touch dead rodents with bare hands, use gloves or use tools such as tongs when disposing them. This product contains a bittering agent and a dye.

**INSTRUCTIONS FOR SAFE DISPOSAL OF THE PRODUCT AND ITS PACKAGING:** At the end of the treatment, dispose of uneaten bait and the packaging in accordance with local requirements. Use of gloves is recommended. For information on disposal in the UK contact the Environment Agency (<http://www.environment-agency.gov.uk>) or SEPA (<http://www.sepa.org.uk>).

**STORAGE:** Store in a dry, cool and well ventilated place. Keep the container closed and away from direct sunlight. Store in places prevented from the access of children, birds, pets and farm animals.

#### Read attached instructions before use.

To be used only by professional users holding certification demonstrating compliance with UK rodenticide stewardship regime requirements.

May cause damage to organs (blood) through prolonged or repeated exposure. Do not breathe dust. Get medical advice if you feel unwell. Dispose of contents/container to a licensed hazardous-waste disposal contractor or collection site except for empty clean containers which can be disposed of as non-hazardous waste.

#### Active substance:

Brodifacoum 0.029 %

(CAS No.: 56073-10-0)

#### Authorization number:

UK-2015-0857

#### Contains: 3 kg

#### Date of manufacture, date of

#### expiry and batch number:

on the packaging.

WARNING



33062

