

GB - RATIMOR FRESH BAIT

INSTRUCTIONS FOR USE:

Indoor (House mouse (*Mus musculus*), Brown rat (*Rattus norvegicus*)):

Mice: High infestation: Up to 40 g of bait per baiting point spaced 2m apart. Low infestation: Up to 40 g of bait per baiting point spaced 5m apart.

Rats: High infestation: Up to 200 g of bait per baiting point spaced 5m apart. Low infestation: Up to 200 g of bait per baiting point spaced 10m apart.

Application method: Ready-to-use bait to be used in tamper-resistant bait stations. Covered and protected baiting points.

Remove the remaining product at the end of treatment period. Sites under a permanent baiting regime should be inspected regularly in accordance with product label directions. The period between visits should be determined by the technician in charge but will not be longer than every four weeks when permanent baiting is conducted outdoors. For permanent baiting follow any additional instructions provided by the CRRU Guidance on Permanent Baiting. Where possible, prior to the treatment inform any possible bystanders (e.g. users of the treated area and their surroundings) about the rodent control campaign. Consider preventive control measures (e.g. plug holes, remove potential food and drinking as far as possible) to improve product intake and reduce the likelihood of reinfestation. To reduce risk of secondary poisoning, search for and remove dead rodents during treatment at frequent intervals, in line with the recommendations provided by the relevant code of best practice. Do not use this product in pulsed baiting treatments. Permanent baiting is strictly limited to sites with a high potential for reinfestation when other methods of control have proven insufficient. The permanent baiting strategy shall be periodically reviewed in the context of integrated pest management (IPM) and the assessment of the risk for re-infestation. When placing bait points close to water drainage systems, ensure that bait contact with water is avoided.

Outdoor around buildings (House mouse (*Mus musculus*), Brown rat (*Rattus norvegicus*)):

Mice: High infestation: Up to 40 g of bait per baiting point spaced 2m apart. Low infestation: Up to 40 g of bait per baiting point spaced 5m apart.

Rats: High infestation: Up to 200 g of bait per baiting point spaced 5m apart. Low infestation: Up to 200 g of bait per baiting point spaced 10m apart. Burrow baiting: 10 – 200 g of bait per burrow.

Application method: Ready-to-use bait to be used in tamper-resistant bait stations. Covered and protected baiting points. Direct application of ready-to-use bait into the burrow.

Protect bait from the atmospheric conditions. Place the baiting points in areas non-labile to flooding. Replace any bait in baiting points in which bait has been damaged by water or contaminated by dirt. Remove the remaining product at the end of treatment period (except when directly applied into burrows). Bait must be placed to minimise the exposure to non-target species and children. Cover or block the entrances of baited burrows to reduce the risks of bait being rejected and spilled. Follow any additional instructions provided by the CRRU UK Code of Best Practice (or equivalent). For outdoor use, baiting points must be covered and placed in strategic sites to minimise the exposure to non-target species. Sites under a permanent baiting regime should be inspected regularly in accordance with product label directions. The period between visits should be determined by the technician in charge but will not be longer than every four weeks when permanent baiting is conducted outdoors. For permanent baiting follow any additional instructions provided by the CRRU Guidance on Permanent Baiting. Where possible, prior to the treatment inform any possible bystanders (e.g. users of the treated area and their surroundings) about the rodent control campaign. Consider preventive control measures (plug holes, remove potential food and drinking as far as possible) to improve product intake and reduce the likelihood of reinfestation. To reduce risk of secondary poisoning, search for and remove dead rodents during treatment at frequent intervals, in line with the recommendations provided by the relevant code of best practice. Do not use this product in pulsed baiting treatments. Permanent baiting is strictly limited to sites with a high potential for reinfestation when other methods of control have proven insufficient. The permanent baiting strategy shall be periodically reviewed in the context of integrated pest management (IPM) and the assessment of the risk for re-infestation. When placing bait points close to surface waters (e.g. rivers, ponds, water channels, dykes, irrigation ditches) or water drainage systems, ensure that bait contact with water is avoided.

Outdoor open areas and waste dumps (Brown rat (*Rattus norvegicus*)):

High infestation: Up to 200 g of bait per baiting point spaced 5m apart. Low infestation: Up to 200 g of bait per baiting point spaced 10m apart. Burrow baiting: 10 – 200 g of bait per burrow.

Application method: Ready-to-use bait to be used in tamper-resistant bait stations. Covered and protected baiting points. Direct application of ready-to-use bait into the burrow.

Protect bait from the atmospheric conditions. Place the bait stations in areas non-labile to flooding.

UNICHEM

Manufacturer and approval holder:
UNICHEM d.o.o., Sinja Gorica 2, 1360 Vrhnika,
Slovenia, Tel.: +386 1 7558 152,
unichem@unichem.si, www.unichem.si

Distributor:

Killgerm Chemicals Ltd, P.O. BOX 2,
Ossett, West Yorkshire, WFS 9NA, UK,
Tel.: 01924 268 400, Fax: 01924 264 757,
http://www.killgerm.com



WITH
SPECIAL
AROMA

RATIMOR®

BROMADIOLONE 0,05 g/kg

Fresh Bait
80 g



Professional
Solutions

4 kg

Replace any bait in baiting points in which bait has been damaged by water or contaminated by dirt. Remove the remaining product at the end of treatment period (except when directly applied into burrows). Bait must be placed to minimise the exposure to non-target species and children. Cover or block the entrances of baited burrows to reduce the risks of bait being rejected and spilled. Follow any additional instructions provided by the CRRU UK Code of Best Practice (or equivalent). Sites under a permanent baiting regime should be inspected regularly in accordance with product label directions. The period between visits should be determined by the technician in charge but will not be longer than every four weeks when permanent baiting is conducted outdoors. For permanent baiting follow any additional instructions provided by the CRRU Guidance on Permanent Baiting. For outdoor use, baiting points must be covered and placed in strategic sites to minimise the exposure to non-target species. Where possible, prior to the treatment inform any possible bystanders (e.g. users of the treated area and their surroundings) about the rodent control campaign. To reduce risk of secondary poisoning, search for and remove dead rodents during treatment at frequent intervals, in line with the recommendations provided by the relevant code of best practice. Permanent baiting is strictly limited to sites with a high potential for reinfestation when other methods of control have proven insufficient. The permanent baiting strategy shall be periodically reviewed in the context of integrated pest management (IPM) and the assessment of the risk for re-infestation. Do not use this product in pulsed baiting treatments. When placing bait points close to surface waters (e.g. rivers, ponds, water channels, dykes, irrigation ditches) or water drainage systems, ensure that bait contact with water is avoided.

Sewers (Brown rat (*Rattus norvegicus*)):

High infestation: 200 g of bait per manhole. Low infestation: Up to 200 g of bait per manhole.

Application method: Ready-to-use bait to be anchored or applied in bait stations preventing the bait from getting into contact with waste water. Covered and protected baiting points.

Bait must be applied in a way so that they do not come into contact with water and are not washed away. Where possible, it is recommended that the treated area is revisited every 4 weeks at the latest in order to avoid any selection of a resistant population. Do not use this product in pulsed baiting treatments. Permanent baiting is strictly limited to sites with a high potential for reinfestation when other methods of control have proven insufficient. The permanent baiting strategy shall be periodically reviewed in the context of integrated pest management (IPM) and the assessment of the risk for re-infestation.

Read attached instructions before use.

To be used only by professional users holding certification demonstrating compliance with UK rodenticide stewardship regime requirements.

May damage the unborn child. Causes damage to organs (Blood) through prolonged or repeated exposure. Obtain special instructions before use. Do not handle until all safety precautions have been read and understood. Wear protective gloves. Do not eat, drink or smoke when using this product. Wash hands thoroughly after handling. If exposed or concerned: Get medical advice/attention. Get medical advice/attention if you feel unwell. Store locked up. Dispose of contents/container to a licensed hazardous waste disposal contractor or collection site except for empty clean containers which can be disposed of as non-hazardous waste.

Active substance:
Bromadiolone 0.005 %
(CAS No.: 28772-56-7)

Authorization number:
(UK-2012-0656-0001)

Contains: 4 kg
**Date of manufacture, date of expiry
and batch number:**
on the packaging.

DANGER



30740

