

STATUTORY CONDITIONS OF APPROVAL RELATING TO USE:

FOR USE ONLY AS AN INSECTICIDE. FOR USE ONLY BY PROFESSIONAL OPERATORS

Routine Rate: Dilute 25ml in 5L of water and apply at a rate of 50ml per square metre

Single/Initial Rate: Dilute 50ml in 5L of water and apply at a rate of 50ml per square metre

Apply by spray.

Apply at a rate of 50 per metre square. (1L per 20 square metres)

Users must comply in full with all their legal responsibilities under The Control of Pesticide Regulations 1986 (as amended), including complying with all conditions of approval.

Users must comply in full with all their legal responsibilities under of Pesticide Regulations (Northern Ireland) 1987 (as amended), including complying with all conditions of approval.

READ ALL PRECAUTIONS BEFORE USE

HSE 9988-GB, HSE 9988-NI

Killgerm®



Professional insecticides
for professional use

PRECAUTIONS:

TO AVOID RISKS TO MAN AND THE ENVIRONMENT, COMPLY WITH INSTRUCTIONS FOR USE.

- THE COSHH (CONTROL OF SUBSTANCES HAZARDOUS TO HEALTH) REGULATIONS 2002 (AS AMENDED) APPLY TO THE USE OF THIS PRODUCT AT WORK.
- AVOID EXCESSIVE CONTAMINATION OF COVERALLS.
- DO NOT CONTAMINATE FOODSTUFFS, EATING UTENSILS OR FOOD CONTACT SURFACES.
- THIS MATERIAL AND ITS CONTAINER MUST BE DISPOSED OF IN A SAFE WAY.
- DO NOT APPLY DIRECTLY TO LIVESTOCK AND POULTRY.
- EXCLUDE ALL PERSONS AND ANIMALS DURING TREATMENT.
- COVER ALL WATER STORAGE TANKS BEFORE APPLICATION.
- ALL BATS ARE PROTECTED UNDER THE WILDLIFE AND COUNTRYSIDE ACT OF 1981. BEFORE TREATING ANY STRUCTURE USED BY BATS CONSULT NATURAL ENGLAND, SCOTTISH NATURAL HERITAGE OR THE COUNTRYSIDE COUNCIL FOR WALES.
- HAZARDOUS TO BEES.
- DO NOT APPLY TO CLOTHES AND BEDDING.
- WASH HANDS AND EXPOSED SKIN BEFORE MEALS AND AFTER USE.
- KEEP IN A SAFE PLACE.
- DO NOT TOUCH TREATED SURFACES UNTIL DRY.
- COLLECT EGGS AND REMOVE FEED, EXPOSED WATER AND MILK BEFORE APPLICATION.
- NOT TO BE USED ON PET BEDDING, PET SLEEPING QUARTERS OR WHERE PETS FREQUENTLY LIE.
- ACTION SHOULD BE TAKEN TO PREVENT FORAGING BEES GAINING ACCESS TO TREATED NEST. PREFERABLY BY REMOVING THE COMBS OR BLOCKING THE NEST ENTRANCE.
- DO NOT APPLY DIRECTLY TO ANIMALS.

ADDITIONAL INFORMATION FOR NI:

- ALL BATS ARE PROTECTED UNDER THE CONSERVATION (NATURAL HABITATS ETC.) REGULATIONS (NORTHERN IRELAND) 1995. BEFORE TREATING ANY STRUCTURE USED BY BATS, CONSULT NORTHERN IRELAND ENVIRONMENT AGENCY.

Marketing Company:
Killgerm Chemicals Ltd, Wakefield Road, Osselt, West Yorkshire, WF5 9AJ
t 01924 268400 e sales@killgerm.com www.killgerm.com

HAZARD PHRASES:

HARMFUL IF SWALLOWED.
CAUSES SEVERE EYE DAMAGE.
MAY CAUSE DAMAGE TO ORGANS THROUGH PROLONGED OR REPEATED EXPOSURE.
VERY TOXIC TO AQUATIC LIFE WITH LONG LASTING EFFECTS.

PRECAUTIONARY PHRASES:

DO NOT BREATHE VAPORS
WASH WITH SOAP AND WATER THOROUGHLY AFTER HANDLING.
DO NOT EAT, DRINK OR SMOKE WHEN USING THIS PRODUCT.
AVOID RELEASE TO THE ENVIRONMENT
WEAR PROTECTIVE GLOVES/PROTECTIVE CLOTHING/EYE OR FACE PROTECTION.
IF SWALLOWED: CALL A POISON CENTER OR DOCTOR/PHYSICIAN.
RINSE MOUTH.
IF IN EYES: RINSE CONTINUOUSLY WITH WATER FOR SEVERAL MINUTES. REMOVE CONTACT LENSES, IF PRESENT AND EASY TO DO SO. CONTINUE RINSING.
GET MEDICAL ADVICE/ATTENTION IF YOU FEEL UNWELL.
IMMEDIATELY CALL A POISON CENTRE OR DOCTOR/PHYSICIAN.
COLLECT SPILLAGE
DISPOSE OF CONTENTS AND CONTAINER TO HAZARDOUS OR SPECIAL WASTE COLLECTION POINT, IN ACCORDANCE WITH LOCAL, REGIONAL, NATIONAL AND/OR INTERNATIONAL REGULATION.
SAFETY DATA SHEET AVAILABLE ON REQUEST
SHELF LIFE: USE WITHIN 2 YEARS OF MANUFACTURE. PROTECT FROM FROST.



ANGER

UFI: TH00-60WE-800J-TT5F

Contains: Cypermethrin 10.0% + Tetramethrin 0.9%

Vazor®
PYRETHROID BASED INSECTICIDE CONCENTRATE
CYPERMAX PLUS

CYPERMAX PLUS

PYRETHROID BASED INSECTICIDE CONCENTRATE

Insecticide for professional use against crawling and flying insects such as: Flies, Moths, Beetles, Fleas, Ticks, Bedbugs, Cockroaches, crickets, silverfish, bedbugs, booklice, mites, ants, mosquitoes, wasps and centipedes.

For use in and around domestic properties, public and municipal areas, refuse tips, poultry houses, slaughter houses and manure heaps.

Applications:

Remove cap from dosing chamber on bottle. Squeeze bottle to fill dosing chamber. Add to 5L of clean water and mix well.

Routine Rate: Dilute 25ml in 5L of water and apply at a rate of 50ml per square metre

Single/Initial Rate: Dilute 50ml in 5L of water and apply at a rate of 50ml per square metre

Apply by spray.

Apply at a rate of 50ml per metre square. (1L per 20 square metres)

CONTAINS: CYPERMETHRIN 10.0% + TETRAMETHRIN 0.9%

Contents: Up to 5L

Batch number-see separate sticker
D.O.M see separate sticker

LABEL REF-AUG26



USE BIOCIDES SAFELY
Always read the label and
product information before use